

# Hepatitis B virus infection amongst pregnant women in North-Eastern Nigeria- A call for action

AB Olokoba, FK Salawu<sup>1</sup>, A Danburam<sup>1</sup>, LB Olokoba<sup>2</sup>, JK Midala<sup>1</sup>, LH Badung<sup>1</sup>, AWO Olatinwo<sup>3</sup>

Department of Medicine, <sup>2</sup>Ophthalmology and <sup>3</sup>Obstetrics and Gynaecology, University of Ilorin Teaching Hospital, Ilorin, <sup>1</sup>Federal Medical Centre, Yola, Nigeria

## Abstract

**Background:** It is well known that Hepatitis B virus infection is endemic in Nigeria. Even though studies have been carried out on Hepatitis B virus infection in different parts of Nigeria, and in different sub-groups of individuals, information regarding the prevalence of Hepatitis B virus infection in pregnant women is scanty especially from the North-eastern region of Nigeria. We therefore determined the seroprevalence of Hepatitis B surface antigen (HBsAg) amongst pregnant women in North Eastern Nigeria.

**Materials and Methods:** A hospital-based cross-sectional study was carried out. The setting was the ante-natal clinic of the Federal Medical Centre, Yola, Nigeria. The duration of the study was from July 2008 to December 2008. Two hundred and thirty-one consecutively recruited pregnant women were screened for Hepatitis B surface antigen. Positive samples were re-tested using ELISA to eliminate false positives. Their biodata were obtained using a questionnaire to establish the presence of possible risk factors such as blood transfusion, surgery, etc. Written informed consent was obtained from each woman.

**Results:** Out of the 231 pregnant women tested, nineteen of them were seropositive for Hepatitis B virus infection giving an infection rate of 8.2%. Women in the age group 25-29 years had the highest HBV infection rate.

**Conclusion:** This study confirms a high seroprevalence of Hepatitis B virus infection amongst pregnant women. It is recommended that pregnant women should be routinely screened for Hepatitis B virus infection as part of antenatal care services.

**Key words:** Hepatitis B virus, Nigeria, pregnancy, seroprevalence

**Date of Acceptance:** 21-Apr-2010

## Introduction

Hepatitis B virus (HBV) is a DNA virus causing hepatitis in humans. It accounts for 400 million chronic infections worldwide,<sup>[1]</sup> and is hyperendemic in sub-Saharan Africa and Asia.<sup>[2,3]</sup> It is thought to be the main aetiological factor in over 75% of chronic liver diseases.<sup>[3]</sup> Transmission of HBV results from exposure to infectious blood or body fluids, unprotected sexual contact, blood transfusion, reuse of contaminated needles and syringes, and vertical transmission from mother to child.<sup>[4,5]</sup> Other high-risk adult populations include persons with multiple heterosexual partners, men who have sex with men and healthcare

workers. Without intervention, the risk of peri-natal HBV transmission is greatest for infants born to women who are HBeAg-positive, with infectivity rate of 70% to 90% at 6 months of age, and about 90% of these children remain chronically infected.<sup>[6]</sup> The risk of peri-natal infection among infants born to HBeAg-negative mothers ranges from 10% to 40%, with 40–70% of these infected infants remaining chronically infected. Children born to HBsAg-positive mothers who do not become infected during the peri-natal period remain at a high risk of infection during early childhood. HBV-related end-stage liver disease or

### Address for correspondence:

Dr. Olokoba AB  
Gastroenterology Unit, Department of Medicine, University of Ilorin Teaching Hospital, P.M.B 1459, Ilorin, Kwara State, Nigeria.  
E-mail: drabolokoba@yahoo.com

### Access this article online

Quick Response Code:



Website: [www.njcponline.com](http://www.njcponline.com)

DOI: 10.4103/1119-3077.79232

PMID: 21493984

hepatocellular carcinoma (HCC) are responsible for over 1 million deaths per year and currently represent 5–10% of cases of liver transplantation.<sup>[7-9]</sup> HCC is one of the most common cancers worldwide and HBV is responsible for at least 75% of this cancer.

HBV can also be transmitted between family members within households possibly by contact of non-intact skin or mucous membrane with secretions or saliva containing HBV.<sup>[4,5]</sup>

Testing for HBV infection in pregnancy is important in view of the morbidity and mortality of the host (pregnant women), its effect on the process of parturition, and the risk of vertical transmission from mother-to-child.<sup>[10]</sup> The contaminated newborn most often remains a chronic carrier with the attendant consequences of liver cirrhosis, and HCC. Mother-to-child transmission can be avoided by vaccination of the newborn.<sup>[10]</sup> This intervention to stop vertical transmission can only be applied when the status of the pregnant woman is known. Although studies have been carried out on HBV infection in different regions of Nigeria in different sub-groups of individuals like blood donors, information regarding HBV infection in pregnancy in Nigeria is sparse especially from the North-eastern region

even though Nigeria falls within a high endemic region as regards the prevalence of HBV infection. We, therefore, investigated the occurrence of HBV infection amongst pregnant women in North-eastern Nigeria.

## Materials and Methods

The serum samples (about 2.5 ml) of 231 consecutively recruited pregnant women attending the ante-natal clinic were screened for HBV using latex rapid agglutination slide test to detect hepatitis B surface antigen (HBsAg). Reactive samples were further confirmed for HBsAg using enzyme-linked immunosorbent assay (ELISA) (Bio Rad, France).

Their biodata were obtained using a structured questionnaire to establish the presence of possible risk factors such as blood transfusion, surgery, etc. Written informed consent was obtained from each woman.

The study was approved by the Ethics and Research committee of the Federal Medical Centre, Yola, Nigeria.

## Analysis

The data obtained were analysed using the statistical package for social sciences (SPSS, version 10.0) statistical software.

## Results

At the conclusion of the study, 231 pregnant women were screened. Nineteen of them were seropositive for HBV infection.

### Age

The age of the women studied ranged from 17 to 44 years with a mean of  $27.8 \pm 5.2$  years. There was a steady increase in the age groups of the women, with a peak in the 25–29 year age group and a decline towards the 40–44 year age group. Majority of the women were in the age group 25–29 years (i.e., 39.8%). Women in the age group 25–29 years had the highest HBV infection rate Table 1.

### Level of education

Majority of the women tested (51.5%) were educated up to the tertiary level, while 2.6% had no form of education. Table 2.

### History of blood transfusion and surgery

Majority of the women tested (93.5%) never had blood

**Table 1: HBsAg Seroprevalence rate among various age groups**

Age groups (Years)	Frequency	Percentage	HBV (n)(%)
15–19	10	4.3	1 (0.4)
20–24	49	21.2	4 (1.7)
25–29	92	39.8	5 (2.2)
30–34	51	22.1	3 (1.3)
35–39	24	10.4	4 (1.7)
40–44	5	2.2	2 (0.9)
Total	231	100	19 (8.2)

**Table 2: Level of education**

Level of education	Frequency	Percentage
None	6	2.6
Quranic	16	6.9
Primary	19	8.2
Secondary	71	30.7
Tertiary	119	51.5
Total	231	100.0

**Table 3: History of blood transfusion and surgery**

Blood transfusion	Frequency	Percentage	Surgery	Frequency	Percentage
No	216	93.5	No	195	84.4
Yes	15	6.5	Yes	36	15.6
Total	231	100.0		231	100.0

transfusion and majority of them (84.4%) also never had surgery Table 3.

## Discussion

Sexually transmitted infections (STIs) and HIV/AIDS are widespread in the developing countries and constitute a major public health problem in sub-Saharan Africa.<sup>[11,12]</sup> The classification of high endemicity for HBV infection has been defined as HBsAg greater than 7% in an adult population.<sup>[13]</sup> The HBsAg seropositivity of 8.2% among pregnant women in our study shows that the North-eastern region like other parts of Nigeria is endemic for HBV infection. This result is in conformity with an earlier finding that sub-Saharan Africa has HBV carrier rate ranging from 9% to 12%.<sup>[14]</sup>

From this study, a seroprevalence rate of 8.2% was found for HBV infection in pregnant women in North-eastern Nigeria. This figure is higher than the 2.9% found in pregnant women in Port Harcourt, South-south Nigeria by Obi et al.<sup>[15]</sup> It is also higher than the 6.2% found in pregnant women in Sierra Leone by Wurie et al.,<sup>[16]</sup> the 2.5% found by Sahaf et al.<sup>[17]</sup> in pregnant Iranian women, and the 1.53% found by Todd et al.<sup>[18]</sup> amongst pregnant Afghan women attending government maternity hospitals in Kabul.

The figure of 8.2% from this study, is however lower than the 11.0% found by Mbaawuaga et al.<sup>[19]</sup> among the pregnant women in Makurdi, North-central Nigeria. It is also lower than the 11.6% found by Harry et al.,<sup>[20]</sup> and the 12.6% found by Jombo et al.<sup>[21]</sup> among pregnant women in Maiduguri, North-eastern Nigeria, and a rural community in North-central Nigeria, respectively. Furthermore, it is also lower than the 13.8% found by Roingard et al.<sup>[22]</sup> in pregnant Senegalese women in Dakar. Lastly, it is also lower than the 63.3% found by Imade et al.<sup>[23]</sup> in Jos, North-central Nigeria amongst pregnant Nigerian women. The wide variations in the seroprevalence of HBV in pregnant women from the literature may be due to geographical variation, differences in cultural practices, sexual behaviour and practices, and differences in the test methods employed to detect HBV infection. Imade et al.<sup>[18]</sup> determined the seroprevalence of HBV infection by the detection of hepatitis B virus core antigen (HBcAg) in Jos, whereas our study employed the use of HBsAg. Most of the studies cited from the literature did not distinguish between recent and past HBV infection. We did not assay for other serological markers of HBV infection in our study such as anti-HBs and anti-HBc antibodies, which are indicators of previous exposure to HBV infection. If these markers were assayed for, the actual seroprevalence rate would most probably be much higher than the present reported figures. Screening for HBsAg alone does not fully reflect the epidemiology of the disease as it could indicate a carrier state, viral replication, or chronic hepatitis. Therefore, our study did not differentiate carriers of HBsAg from those with active

infection. A high frequency of HBV infection was found in the 25–29 years age group followed by the 20–24 and the 30–34 years age groups. This age group of 25–34 years is the most sexually active and fertile. This may explain the high prevalence of HBV infection in them. Majority of the pregnant women tested had tertiary education. This may be because this study was hospital-based in an urban centre. The implication of HBV infection in pregnancy is the risk of vertical transmission from mother-to-child. The contaminated newborn most often remains a chronic carrier with the attendant consequences of liver cirrhosis, and HCC. Mother-to-child transmission can be avoided by vaccination of the newborn. This intervention to stop vertical transmission can only be applied when the status of the pregnant woman is known. There is an urgent need to vaccinate all infants born to mothers who are carriers of HBsAg.

## Conclusion

This study confirms a high seroprevalence of HBV infection amongst pregnant women in North-eastern Nigeria. It is recommended that pregnant women should be routinely screened for HBV infection as part of antenatal care services.

## References

1. Alter MJ. Epidemiology of viral hepatitis and HIV co-infection. *J Hepatol* 2006;44:S6-9.
2. Gashau W, Mohammed I. Hepatitis B viral markers in Nigerian patients with primary liver carcinoma. *Trop Geogr Med* 1991;43:64-7.
3. Isselbacher KJ, Wands JR. Neoplasms of the Liver. In: Wilson JD, Braunwald E, Isselbacher K, (eds). *Harrison's principles of Internal Medicine*. 12<sup>th</sup> ed. New York: Mc Graw Hill; 1991. p. 1350-2.
4. Petersen NJ, Barrett DH, Bond WW, Berquist KR, Farew MS, Bender TR, et al. Hepatitis B surface antigen in saliva, impetiginous lesions, and the environment in two remote Alaskan villages. *Appl Environ Microbiol* 1976;32:572-4.
5. Available from: <http://www.health.vic.gov.au/ideas/diseases/hepb> [Last accessed on 2010 Dec 17].
6. Mc Mahon BJ, Alward WL, Hall DB. Acute hepatitis B virus infection: relation of age to the clinical expression of disease and subsequent development of the carrier state. *J Infect Dis* 1985;151:599-603.
7. European association for the study of the liver, International Consensus Conference on Hepatitis B. Consensus statement. *J Hepatol* 2003;39:S3-25.
8. Ganem D, Prince AM. Hepatitis B virus infection natural history and clinical consequences. *N Engl J Med* 2004;350:1118-29.
9. Maddrey WC. Hepatitis B: an important public health issue. *J Med Virol* 2000;61:362-6.
10. Kane M. Global programme for control of hepatitis B infection. *Vaccine* 1995;13:S47-9.
11. Osmond M, Dennis H. Classification and staging of HIV disease. In: Cohen PT, Sande MA, Volberding PA, (eds). *The AIDS knowledge base*. 2<sup>nd</sup> ed. New York: Little Brown; 1994;312-20.
12. Nwokedi EE, Iiyasu Z, Dikko AU, Azeze AO, Mohammed B. Syphilis in a Nigerian paramilitary agency: a need for treatment policy. *Ann Afr Med* 2005;4:177-9.
13. Uneke CJ, Ogbu O, Inyama PU, Anyanwu GI, Njoku MO, Idoko JH. Prevalence of hepatitis-B surface antigen among blood donors and human immunodeficiency virus-infected patients in Jos, Nigeria. *Mem Inst Oswaldo Cruz* 2005;100:13-6.
14. Kiire CF. The epidemiology and Prophylaxis of hepatitis B in Sub-Saharan Africa: A review from tropical and subtropical Africa. *Gut* 1996;38:5-12.
15. Obi RK, Umeh SC, Okurede OH, Iroagba II. Prevalence of hepatitis B virus infection among pregnant women in an antenatal clinic in Port Harcourt,

- Nigeria. *Afr J Clin Exp Micro* 2006;7:78-82.
16. Wurie IM, Wurie AT, Gevao SM. Seroprevalence of hepatitis B virus among middle to high socio-economic antenatal population in Sierra Leone. *West Afr J Med* 2005;24:18-20.
  17. Sahaf F, Tanomand A, Montazam H, Sany AA. Seroprevalence of Hepatitis C, Hepatitis B and HIV and co-infection among pregnant women: a retrospective study in 2006 at Malekan city, Iran. *Res J Med Sci* 2007;1:138-41.
  18. Todd CS, Ahmadzai M, Atiqzai F, Miller S, Smith JM, Ghazan SA, *et al.* Seroprevalence and correlates of HIV, Syphilis, and hepatitis B and C virus among intrapartum patients in Kabul, Afghanistan. *BMC Infect Dis* 2008;8:119.
  19. Mbaawuaga EM, Ebenebeaku MNO, Okopi JA, Damen JG. Hepatitis B virus (HBV) infection among pregnant women in Makurdi, Nigeria. *Afr J Biochem Res* 2008;11:155-9.
  20. Harry TO, Bajani M D, Moses AE. Hepatitis B Virus infection among blood donors and pregnant women in Maiduguri, Nigeria. *East Afr Med J* 1994;71: 596-7.
  21. Jombo GTA, Egah DZ, Banwat EB. Hepatitis B Virus infection in a rural Settlement of Northern Nigeria. *Niger J Med* 2005;14:425-8.
  22. Roingear P, Diouf A, Sankale JL, Boye C, Mboup S, Diadhiou F *et al.* Perinatal transmission of hepatitis B virus infection in Senegal, West Africa. *Viral Immunol* 1993;6:65-73.
  23. Imade GE, Sagay AS, Ugwu BT, Thacher TD, Ford RW. Seroprevalence of Hepatitis B and HIV infections in pregnant women in Nigeria. *J Med Trop* 2004;6:15-21.

Source of Support: Nil, Conflict of Interest: Nil.



#### Announcement

### “QUICK RESPONSE CODE” LINK FOR FULL TEXT ARTICLES

The journal issue has a unique new feature for reaching to the journal's website without typing a single letter. Each article on its first page has a “Quick Response Code”. Using any mobile or other hand-held device with camera and GPRS/other internet source, one can reach to the full text of that particular article on the journal's website. Start a QR-code reading software (see list of free applications from <http://tinyurl.com/yzlh2tc>) and point the camera to the QR-code printed in the journal. It will automatically take you to the HTML full text of that article. One can also use a desktop or laptop with web camera for similar functionality. See <http://tinyurl.com/2bw7fn3> or <http://tinyurl.com/3ysr3me> for the free applications.